

Pharmacoepidemiological study of prescription pattern of analgesics, antipyretics, and nonsteroidal anti-inflammatory drugs at a tertiary health care center

Mohammed A. Al-Homrany, MD, FRCPC, Yacoub M. Irsheid, MD, PhD.

ABSTRACT

Objectives: To examine the outpatients' prescribing pattern of doctors to analgesics, antipyretics, and nonsteroidal anti-inflammatory drugs (NSAIDs) at a tertiary care setting in Southwestern Saudi Arabia.

Methods: A retrospective review of one-year outpatient prescriptions kept by the pharmacy department at Aseer Central Hospital during the period 8 April, 2000 until 7 April, 2001. Five working days per week of each season were sampled randomly and systemically as every other prescription. The type and number of drugs prescribed, patient's diagnoses and age, and the prescribing physician were analyzed. The study sample included 3796 prescriptions.

Results: There was no significant seasonal variation in the pattern of prescription. The most commonly prescribed agent was paracetamol followed in decreasing frequency by ibuprofen, diclofenac, and aspirin. In few of the prescriptions combination of analgesics, antipyretics, and NSAIDs were used. One fourth of prescriptions, the diagnosis was not mentioned or was not clearly written, infection in 40%, and musculoskeletal disorders in 17.7%. The rest were a variety of problems for some of which, the drugs were used inappropriately. Aspirin was used exclusively in adults for cardio-protection, while paracetamol was used mainly as analgesic-antipyretic over all age groups

Conclusions: Some of prescriptions suffered from significant deficiencies. In light of the serious adverse effects of analgesics, antipyretics and NSAIDs, education of physicians on rational use of such drugs, and prescription writing seems necessary.

Saudi Med J 2007; Vol. 28 (3): 369-374

From the Departments of Internal Medicine (Al-Homrany) and the Department of Clinical Pharmacology (Irsheid), College of Medicine, King Khalid University, PO Box 641, Abha, Kingdom of Saudi Arabia.

Received 30th April 2006. Accepted 12th November 2006.

Address correspondence and reprint request to: Dr. Prof. Mohammed A. Al-Homrany, Department of Internal Medicine, College of Medicine, King Khalid University, PO Box 641, Abha, Kingdom of Saudi Arabia. Fax. +966 (7) 2313165. E-mail: mhomrany@yahoo.com

Analgesics, antipyretics, and nonsteroidal anti-inflammatory drugs (NSAIDs) are among the most commonly prescribed drugs in clinical practice.¹⁻³ They are commonly used for inflammatory disorders of the musculoskeletal system. They constitute a heterogeneous group of compounds with the common ability to inhibit cyclooxygenase, and thus, prostaglandin synthesis.^{4,5} They also have analgesic and antipyretic actions. These drugs have a wide range of adverse effects, including but not limited to, gastric ulceration, gastrointestinal bleeding, inhibition of platelet function, inhibition of induction of labor, renal dysfunction, sodium and water retention, and hypersensitivity reactions.^{4,5} Paracetamol is a non-opioid analgesic that have antipyretic activity, however, it is not useful as inflammatory agent and lacks many of the adverse effects of NSAIDs. It is thought to inhibit cyclooxygenase in the central nervous system (CNS). It is much better tolerated than the NSAIDs as analgesic-antipyretic. Only in overdose, it can produce hepatic damage. In few cases it has produced renal injury.^{4,5} However, these drugs are perceived by patients as effective for the conditions for which they are prescribed.⁶⁻⁸ Paracetamol is generally considered to be safe analgesic-antipyretic when used at therapeutic doses. Recent evidence has suggested that it can exacerbate bronchial asthma and chronic obstructive pulmonary disease (COPD) and reduce lung function.⁹⁻¹² The prevalence of end-stage renal disease was higher in Aseer, Southwest region of Saudi Arabia. In many cases, the diagnosis was not found.¹³ The authors proposed that the use of analgesics, antipyretics, and NSAIDs might be a contributing factor, especially when used for a long time and at a higher cumulative dose, since analgesics, antipyretics, and NSAIDs are available over-the-counter in Saudi Arabia.

Many studies have shown that NSAIDs are overused and some times irrationally prescribed, and that physicians should be educated to practice evidence-based medicine in their use. Vlahovic-

Palcevski et al¹⁴ reported that evidence-based medicine was not the leading factor in prescribing NSAIDs during 2000 in Rijeka, Croatia, and Stockholm, Sweden. Bernal-Delgado et al¹⁵ have reported that evidence-based outreach visits have improved the physician's behavior on NSAID prescription. Paul and Chauhan¹⁶ reported on the significant use of aspirin in patients with bronchial asthma, where this drug is contraindicated and concluded that awareness programs on rational prescribing of NSAIDs in India are needed.

Although the prescription format may vary slightly from one country to another, most agree on the core elements that should be included in the prescription order.¹⁷⁻²⁰ These essential elements are: name, address, and telephone number of the prescriber, name and address of the patient, patient's age and weight particularly for the extremes of age, date of prescription, name of drug (preferably the generic name), strength and dosage form, dose and frequency of administration, quantity prescribed, the reason for prescribing the drug, instructions for use, and the signature of the prescriber. In Saudi Arabia, all of these requirements are recommended and are available in local regulations. In addition, the physician is required to stamp the prescription. The stamp usually contains name, title, and address of physician.

This work was designed to examine the pattern of prescriptions of analgesics, antipyretics, and NSAIDs in a tertiary health care setting located within the southwestern part of Saudi Arabia, and to identify possible deficiencies in indications for use, patients' age and inappropriate medical indications, and provide suggestions for a more rational prescription behavior for such drugs.

Methods. Aseer Central Hospital is located in Abha City and is the only referral hospital at the tertiary care level in Aseer region. It is utilized by the College of Medicine, King Khalid University for training of Medical Students.

All outpatient prescriptions from within the hospital, irrespective of the clinic of origin, received by and kept in the pharmacy were the target of the study. One-year prescriptions during the period 8 April 2000 until 7 April 2001 were analyzed retrospectively. This period was divided into 4 seasons: spring, summer, fall, and winter. One week (5 working days) of each season was sampled randomly and systemically as every other prescription. Of these, all prescriptions containing paracetamol and NSAIDs were analyzed for the type and number of drugs prescribed, the diagnosis for which they were given, the age of the patient and the prescribing physician. The local "Ethics Committee" at Aseer Central Hospital approved the study. The physicians involved were not

aware of the conduction of the study. The data generated were fed to the Statistical Package for Social Sciences program and simple descriptive statistics were used to analyze results.

Results. Out of 3796 prescriptions reviewed, 2081 (54.8%) of which contained paracetamol or NSAIDs. Paracetamol was the most commonly prescribed among the analgesics, antipyretics, and NSAIDs (63.9%). The next most often prescribed analgesics, antipyretics, and NSAID was ibuprofen, which was present in 14.9% of such prescriptions followed by diclofenac (9%) and aspirin (6.7%) (Table 1). Indomethacin was less frequently prescribed (1.6%). Combinations of paracetamol with NSAIDs were present in 3% of prescriptions and NSAID combinations (2 or 3) in 0.9% of prescriptions. There was no significant seasonal variation in the prescribing frequency of the most often prescribed analgesics, antipyretics, and NSAIDs. Using the analysis of variance to compare the frequency of NSAID use with the season, the F statistic was 0.658 and the *p*-value was 0.578. However, paracetamol was slightly less prescribed in spring (19.5%), while ibuprofen was slightly more prescribed in fall (33.1%). Diclofenac was more prescribed in winter (36.9%) and less so in spring and fall (17.1% each). Aspirin was uniformly prescribed over the seasons. The prescribing physician was not mentioned in 80.3% of prescriptions, while emergency room doctors, general practitioners, and specialists were responsible for 4.2%, 8.9% and 6.6% of prescriptions, respectively.

The distribution of drugs prescribed according to the diagnosis was presented in Table 2. Paracetamol was most often prescribed for infections (59%), followed by musculoskeletal pain or trauma, in addition to headache and fever (9.8%), apparently as an analgesic-antipyretic. In 19.8% of the prescriptions for paracetamol, the indication was either not mentioned or not clearly written and in 1.9% of prescriptions the diagnosis written in the prescriptions was either bronchial asthma or COPD.

Aspirin was found in 6.7% of prescriptions containing analgesics, antipyretics, and NSAIDs. In 36.4% of prescriptions containing aspirin, the indication was cardiovascular disorders. Only in 4 prescriptions aspirin was prescribed for transient ischemic attacks, while in 46.3% the indication was not mentioned or not clearly written. However, the dose in all of the cases was 325 mg apparently; it was prescribed as an antiplatelet agent.

Ibuprofen was found in 14.9% of prescriptions containing analgesics, antipyretics, and NSAIDs (Table 1). In 44.9% of the prescriptions the indication was musculoskeletal pain or trauma, in 37.6% of the prescriptions the indication was either not mentioned

or not clearly written, in 5.7% it was infections and in 2.2% of prescriptions containing ibuprofen, the diagnosis mentioned in the prescription was gastritis and gastroenteritis.

Diclofenac was found in 9% of prescriptions containing analgesics, antipyretics, and NSAIDs (Table 1). In 50.5% of prescriptions containing diclofenac, the indication was musculoskeletal pain or trauma, while in 21.9% of prescriptions the diagnosis was not mentioned or not clearly written. In 6.8% of prescriptions the indication was renal colic and 9.4% it was infections.

Indomethacin was found in 1.6% of prescriptions containing analgesics, antipyretics and NSAIDs (Table 1). In 34.3% of prescriptions containing indomethacin, the indication was musculoskeletal pain or trauma,

while in 31.4% the indication was not mentioned or not written clearly. In 14.3% of such prescriptions the indication was renal colic and in 8.6% it was nonspecific abdominal pain.

A combination of paracetamol with one or 2 NSAIDs was found in 3% of prescriptions containing analgesics, antipyretics, and NSAIDs (Table 1). In 50.8% of these the indication was musculoskeletal disorders, while in 20% the indication was not mentioned or not clearly written. In 15.4% the indication was infections. Combination of 2 or 3 NSAIDs was present in 0.9% of prescriptions containing analgesics, antipyretics, and NSAIDs (Table 1). In 72.2% of these the indication was musculoskeletal disorders and trauma, while in the rest (27.8%) the indication was not mentioned or not written clearly. Arthritis, osteoarthritis, and gout constituted 12% of the musculoskeletal disorders and the rest constituted musculoskeletal pain or trauma.

Concomitant drugs (Table 3) prescribed along with aspirin were mainly (66.8%) those concerning the cardiovascular system. Paracetamol was prescribed along with anti-infective agents (39.7%), antihistamines, decongestants, expectorants or antitussives (32.5%), drugs for gastrointestinal disorders (8.8%), and drugs for bronchial asthma (3.3%). Ibuprofen was mainly prescribed alone (52.7%) and in conjunction with antihistamines, decongestants, expectorants or antitussives (6.5%). Diclofenac was mainly prescribed alone (50.2%) and in conjunction with anti-infective agents (15.5%) and drugs for gastrointestinal disorders (17.8%). Indomethacin was prescribed alone in 24.5%

Table 1 - Distribution of paracetamol and nonsteroidal anti-inflammatory drugs among the prescriptions.

Drug	Number of prescription	Frequency (%)
Paracetamol	1330	(63.9)
Ibuprofen	311	(14.9)
Diclofenac	187	(9)
Aspirin	139	(6.7)
Indomethacin	34	(1.6)
Paracetamol + NSAIDs	62	(3)
NSAID combinations	18	(0.9)
Total	2081	(100)

NSAID - nonsteroidal anti-inflammatory drug

Table 2 - Prescribing frequencies of analgesic, antipyretic and nonsteroidal antiinflammatory drugs according to the diagnosis

Diagnosis	Paracetamol	Ibuprofen	Diclofenac	Aspirin	Indomethacin	Paracetamol + NSAIDs	NSAID combinations	Total
Not mentioned	102 (7.2)	54 (17.2)	20 (10.4)	60 (37)	2 (5.7)	6 (9.2)	2 (11.1)	246
Not clear	179 (12.6)	64 (20.4)	22 (11.5)	15 (9.3)	9 (25.7)	7 (10.8)	3 (16.7)	299
Infections	836 (59)	18 (5.7)	18 (9.4)	1 (0.6)	1 (2.9)	10 (15.4)	0 (0)	884
Musculoskeletal disorders and trauma	91 (6.4)	141 (44.9)	97 (50.5)	4 (2.5)	12 (34.3)	33 (50.8)	13 (72.2)	391
Renal colic	6 (0.4)	2 (0.6)	13 (6.8)	0 (0)	5 (14.3)	2 (3.1)	0 (0)	28
Abdominal colic & GIT disorders	47 (3.3)	7 (2.2)	6 (3.1)	1 (0.6)	3 (8.6)	1 (1.5)	0 (0)	65
Cardiovascular	11 (0.8)	2 (0.6)	1 (0.5)	59 (36.4)	0 (0)	0 (0)	0 (0)	73
Headache and other pains	48 (3.4)	8 (2.5)	5 (2.6)	3 (1.9)	0 (0)	3 (4.6)	0 (0)	67
Bronchial asthma and COPD	27 (1.9)	0 (0)	0 (0)	1 (0.6)	0 (0)	0 (0)	0 (0)	28
Fever	5 (0.4)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	5
Others	66 (4.7)	18 (5.7)	10 (5.2)	18 (11.1)	3 (8.6)	3 (4.6)	0 (0)	118
Total	1418 (100)	314 (100)	192 (100)	162 (100)	35 (100)	65 (100)	18 (100)	2204*

*The total is more than 2081 as some prescriptions contained two diagnoses, NSAID - Nonsteroidal anti-inflammatory drug, COPD -Chronic obstructive pulmonary disease, GIT -Gastrointestinal tract, GIT disorders - gastritis and gastroenteritis

of prescriptions containing the drug and with drugs for gastrointestinal disorders (38.8%) and anti-infective agents (18.4%). In most cases gastrointestinal drugs prescribed were mainly antacids and anti-ulcer drugs in few cases hyoscine and prokinetic agents.

The age distribution of drugs prescribed is presented in Table 4. The age was mentioned in only 48.9% of the prescriptions. The age was mentioned in 25.9%, 50.7%, 55.3%, 46.5%, and 35.3% of prescriptions containing aspirin, paracetamol, ibuprofen, diclofenac,

and indomethacin. In all prescriptions containing age, aspirin was prescribed exclusively for adults and the elderly, while paracetamol was prescribed for all age groups. The other NSAIDs were mostly prescribed for adults and none for children below one year of age (Table 4).

Discussion. This study describes the pattern of prescription of paracetamol and NSAIDs in outpatient clinics and emergency department of a teaching hospital

Table 3 - Concomitant drugs^a prescribed with analgesic, antipyretic and nonsteroidal anti-inflammatory drugs. Data are number of prescriptions (%)

Concomitant drugs' type	Analgesic, antipyretic or anti-inflammatory drug							Total ^b
	Aspirin	Paracetamol	Ibuprofen	Diclofenac	Indomethacin	Combination		
No other drugs	19 (8.2)	148 (6.6)	186 (52.7)	107 (50.2)	12 (24.5)	48 (55.2)		520
Antiinfective agents	2 (0.9)	893 (39.7)	71 (20.1)	33 (15.5)	9 (18.4)	14 (16.1)		1022
Antihistamines, Decongestants, Expectorants or Antitussives	2 (0.9)	731 (32.5)	23 (6.5)	3 (1.4)	1 (2)	3 (3.5)		763
Drugs for GIT disorders	7 (3)	199 (8.8)	36 (10.2)	38 (17.8)	19 (38.8)	11 (12.6)		310
Drugs for Bronchial Asthma	0 (0)	72 (3.3)	0 (0)	0 (0)	0 (0)	0 (0)		72
Cardiovascular Drugs	155 (66.8)	10 (0.4)	3 (0.9)	4 (1.9)	0 (00)	2 (2.3)		174
Other Drugs	47 (20.2)	196 (8.7)	34 (9.6)	28 (13.2)	8 (16.3)	9 (10.3)		322
Total	232(100)	2249 (100)	353(100)	213(100)	49(100)	87(100)		3183

GIT -Gastrointestinal tract, ^aPrescriptions contained 1-8 drugs,
^bPercentages are not included because this column contain total analgesics, antipyretics or anti-inflammatory drugs

Table 4 - Age-wise prescribing frequencies for analgesics, antipyretics and nonsteroidal anti-inflammatory drugs.

Drug	Number of prescriptions (%)							Total
	Newborn (<2 months)	Infants and toddlers (2 months to <2 years)	Children (2 to <12 years)	Adolescents (12 to <18 years)	Adults (18 to 65 years)	Elderly (>65 years)	Not mentioned	
Aspirin	0 (0)	0 (0)	0 (0)	0 (0)	29 (20.9)	7 (5)	103 (74.1)	139 (100)
Paracetamol	7 (0.5)	107 (8.1)	256 (19.2)	39 (2.9)	255 (19.2)	10 (0.8)	656 (49.3)	1330 (100)
Ibuprofen	0 (0)	1 (0.3)	6 (1.9)	11 (3.5)	149 (47.9)	5 (1.6)	139 (44.7)	311 (100)
Diclofenac	0 (0)	1 (0.5)	3 (1.6)	6 (3.2)	73 (39)	4 (2.1)	100 (53.5)	187 (100)
Indomethacin	0 (0)	0 (0)	1 (2.9)	0 (0)	11 (32.4)	0 (0)	22 (64.7)	34 (100)
Paracetamol + NSAID	0 (0)	0 (0)	4 (6.5)	1 (1.6)	18 (29.0)	0 (0)	39 (62.9)	62 (100)
NSAID combinations	0 (0)	0 (0)	2 (11.1)	2 (11.1)	9 (50.0)	0 (0)	5 (27.8)	18 (100)
Total	7	109	272	59	544	26	1064	2081

NSAIDs - nonsteroidal anti-inflammatory drugs

in Southwestern, Saudi Arabia. Out of the 3796 prescriptions reviewed, 2081 (54.8%) of prescriptions contained analgesics, antipyretics, and NSAIDs. This finding is less than what is reported from primary health care centers in the same area of Saudi Arabia by Mahfouz et al.²¹ They found that the prevalence of prescribing for analgesics-antipyretics (without specifying individual drugs) was 61.9%. Moreover, Ravi Shankar et al.²² from a medical department in a tertiary care hospital in Pokhara, Nepal reported that the frequency of prescribing analgesics was 15.09%, which is much lower than our figure. However, our study was performed on outpatient prescriptions irrespective of the department of origin.

Paracetamol was the most often prescribed drug, apparently as an analgesic-antipyretic, which is considered to be an appropriate indication. It could not be concluded that this drug was over prescribed, while in 11% of prescriptions containing it, no diagnosis was mentioned. Concomitant drugs (antiinfective agents and antihistamines, decongestants, expectorants and antitussives) indicate its use as analgesic-antipyretic. Concerning the NSAIDs, others also have reported that ibuprofen was the most often prescribed NSAID.^{16,23} It is perceived to be the safest NSAID concerning gastrointestinal complications.^{24,25} The use of ibuprofen for gastritis and gastroenteritis could not be justified as the drug is a gastrointestinal irritant and if such patients need an antipyretic, paracetamol should be the first choice. Diclofenac was the second most often prescribed NSAID. Others have reported that diclofenac was the most often prescribe NSAID.¹⁴ The use of diclofenac for infections in conjunction with antiinfective agents and antihistamines, decongestants, expectorants, or antitussives is also inappropriate and considered an overuse of the drug. The rationale for giving indomethacin for nonspecific abdominal pain also could not be understood. The combination of more than one systemic NSAID is also an inappropriate practice, while it may be reasonable to combine a topical agent with a systemic one. It is well-known that the use of NSAIDs is associated with a substantial increase in the risk of gastrointestinal bleeding.^{6,7} In most of the cases, aspirin was appropriately prescribed at low dose as an antiplatelet agent and in combination with drugs used to treat cardiovascular disorders. The use of paracetamol in bronchial asthma and COPD or its co-prescription with drugs for bronchial asthma could not be justified in light of recent evidence that it can reduce lung function and exacerbate bronchial asthma and COPD. Concomitant drugs gave an insight on the indication for the use of NSAIDs. Such agents included mainly antiinfective agents; and antihistamines, decongestants,

expectorants or antitussives; and antacids and anti-ulcer drugs. The last group was given apparently to protect from the gastrointestinal complications of NSAIDs. In some of the prescriptions, NSAIDs were inappropriately prescribed, such as for gastritis and nonspecific abdominal pain, whereas in one fourth of prescriptions, the diagnosis was not mentioned or was not clearly written. One can argue that the prescribing physician was not clear on the indication and tried to avoid writing down the diagnosis, especially when he wrote it unclearly. In such cases, no single letter of the word was clear. Such a practice was performed in a teaching hospital. One should wonder if the same attitude is practiced at a larger scale in primary health care or other health sectors. In such cases, NSAIDs are assumed to have been misused. Knowing that the prevalence of ESRD is high in Saudi Arabia, and in many cases the cause could not be identified and the fact that analgesics, antipyretics, and NSAIDs are available over-the-counter in this country, one can assume an association between these drugs and ESRD. However, such an association remains to be investigated. The age distribution of prescriptions for analgesics, antipyretics, and NSAIDs is noteworthy. Aspirin was prescribed for adults and elderly exclusively, which agrees with the indications for these age groups as a cardioprotective agent.^{26,27} Other NSAIDs were avoided in children below one year of age but mostly prescribed for adults, while paracetamol was prescribed for all age groups.

In conclusion, some of prescriptions for NSAIDs suffered from clear deficiencies in both content and indications. The need for continuing medical education regarding the potential dangers of analgesics, antipyretics, and NSAIDs, the importance of their appropriate and rational utilization and the necessity of appropriate prescription writing regarding both content and indication is obvious.

Acknowledgment. *The authors greatly appreciate the cooperation of the staff of the Pharmacy Department at Aseer Central Hospital during the review of prescriptions, and Mr. Andy Rolex for data entry.*

References

1. Bawazir SA. Prescribing patterns of ambulatory care physicians in Saudi Arabia. *Ann Saudi Med* 1993; 13: 172-177.
2. Yap KB, Chan KM. The prescribing pattern of hospital doctors. *Singapore Med J* 1998; 39: 496-500.
3. Rishi PK, Sangeeta S, Surendra K, Tailang M. Prescription audit: experience in Garhwal (Uttaranchal), India. *Trop Doct* 2003; 33: 76-79.
4. Wagner W, Khanna P, Furst DE. Nonsteroidal anti-inflammatory drugs, disease-modifying antirheumatic drugs, nonopioid analgesics and drugs used in gout. In: Katzung BG, editor. *Basic and Clinical Pharmacology*. 9th ed. International edition, McGraw Hill. 2004. p. 576-603.

5. Roberts II LJ, Morrow JD. Analgesics-antipyretics and anti-inflammatory agents and drugs employed in the treatment of gout. In Hardman JG, Limbird LE, Gilman AG, editors. Goodman and Gilman's the Pharmacological Basis of Therapeutics. 10th ed. International edition, McGraw Hill. 2001. p. 687-731.
6. Lapane KL, Spooner JJ, Mucha L, Straus WL. Effect of nonsteroidal anti-inflammatory drug use on the rate of gastrointestinal hospitalizations among people living in long-term care. *J Am Geriatr Soc* 2001; 49: 577-584.
7. Mellemkjær L, Blot WJ, Sørensen HT, Thomassen L, McLaughlin JK, Nielsen GL, et al. Upper gastrointestinal bleeding among users of NSAIDs: a population-based cohort study in Denmark. *Br J Clin Pharmacol* 2002; 53: 173-181.
8. Hawkey CJ, Cullen DJE, Pearson G, Holmes S, Doherty M, Wilson JV, et al. Pharmacoepidemiology of nonsteroidal anti-inflammatory drug use in Nottingham general practices. *Aliment Pharmacol Ther* 2000; 14: 177-185.
9. Barr RG, Wentowski CC, Curhan GC, Somers SC, Stampfer MJ, Schwartz J, et al. Prospective study of acetaminophen use and newly diagnosed asthma among women. *Am J Respir Crit Care Med* 2004;169: 836-841.
10. Shaheen SO, Newson RB, Henderson AJ, Headley JE, Stratton FD, Jones RW, et al. Prenatal paracetamol exposure and risk of asthma and elevated immunoglobulin E in childhood. *Clin Exp Allergy* 2005; 35: 18-25.
11. McKeever TM, Lewis SA, Smit HA, Burney P, Britton JR, Cassano PA. The association of acetaminophen, aspirin and ibuprofen with respiratory disease and lung function. *Am J Respir Crit Care Med* 2005; 171: 966-971.
12. Karakaya G, Kalyoncu AF. Paracetamol and asthma. *Expert Opin Pharmacother* 2003; 4: 13-21.
13. Al Homrany M, Abualfotoh M. Incidence of treated end-stage renal disease in Aseer region. *Saudi J Kidney Dis Transpl* 1998; 9: 425-429.
14. Vlahovic-Palcevski V, Wettermark B, Bergman U. Quality of nonsteroidal anti-inflammatory drug prescribing in Croatia (Rijeka) and Sweden (Stockholm). *Eur J Clin Pharmacol* 2002; 58: 209-214.
15. Bernal-Delgado E, Galeote-Mayor M, Pradas-Arnal F, Peiro-Moreno S. Evidence-based educational outreach visits: effects on prescriptions of non-steroidal anti-inflammatory drugs. *J Epidemiol Community Health* 2002; 56: 653-658.
16. Paul AD, Chauhan CK. Study of usage pattern of nonsteroidal anti-inflammatory drugs (NSAIDs) among different practice categories in Indian clinical setting. *Eur J Clin Pharmacol* 2005; 60: 889-892.
17. WHO/DAP/94.11. Guide to good prescribing, a practical manual. World health organization, Action programme on Essential Drugs, Geneva, WHO: 1994. p 51-55. Available from URL: http://whqlibdoc.who.int/hq/1994/WHO_DAP_94.11.pdf
18. Lofholm PW, Katzung BG. Rational prescribing and prescription writing. In: Bertram G. Katzung, editor. Basic and Clinical Pharmacology. 9th ed. Chapter 66, International edition, McGraw Hill. 2004. p. 1091-1100.
19. Prescription writing. In: British National Formulary, No 41, British Medical Association and Royal Pharmaceutical Society of Great Britain. 2000. p. 4-5.
20. Anonymous. Safe Writing. In: Lacy CF, Armstrong LL, Goldman MP, Lance LL, editors. "Drug Information Handbook". 8th ed. Lexi Comp. 2000-2001. p. 12.
21. Mahfouz AAR, Shehata AI, Mandil AMA, Al-Erian RAG, Al-Khuzayem A-A M, Kisha A. Prescribing patterns at primary health care level in the Asir region, Saudi Arabia: An epidemiologic study. *Pharmacoepidemiol Drug Saf* 1997; 6: 197-201.
22. Ravi Shankar P, Partha P, Nagesh S. Prescribing patterns in medical outpatients. *Int J Clin Pract* 2002; 56: 549-551.
23. Seager JM, Cullen DJ, Pearson G, Holmes S, Doherty M, Wilson JV, et al. Ibuprofen versus other nonsteroidal anti-inflammatory drugs: use in general practice and patient perception. *Aliment Pharmacol Ther* 2000; 14: 187-191.
24. Henry D, Lim LL, Garcia Rodriguez LA, Perez Gutthann S, Carson JL, Griffin M, et al. Variability in risk of gastrointestinal complications with individual nonsteroidal anti-inflammatory drugs: results of a collaborative meta-analysis. *BMJ* 1996; 312: 1563-1566.
25. Henry D, Dobson A, Turner C. Variability in the risk of major gastrointestinal complications from nonaspirin nonsteroidal anti-inflammatory drugs. *Gastroenterology* 1993; 105: 1078-1088.
26. Masoudi FA, Wolfe P, Havranek EP, Rathore SS, Foody JM, Krumholz HM. Aspirin use in older patients with heart failure and coronary artery disease: national prescription patterns and relationship with outcomes. *J Am Coll Cardiol* 2005; 46: 955-962.
27. Rodondi N, Vittinghoff E, Comuz J, Butler J, Ding J, Satterfield S, et al. Aspirin use for primary prevention of coronary heart disease in older adults. *Am J Med* 2005; 118: 1288.