## Correspondence

# Pregnancy outcome in women with inflammatory bowel disease

To the Editor

Despite the limited number of the studied pregnant with inflammatory bowel diseases (IBD) (n=31), Saleh et al<sup>1</sup> concluded in their study that women with IBD are generally not at great increased risk for maternal, obstetrical, and neonatal complications compared to the general population. This conclusion could be positively viewed in 2 aspects: 1. It is interesting as it precedes a recently published study, which showed that most women with IBD have generally a normal pregnancy outcome, and overall, IBD during pregnancy poses lowto-moderate risks, the main concern being increased risk of impaired fetal growth.2 Moreover, it supports the observation that both disease activity in IBD during pregnancy, and the type of medical treatment do not predispose to abnormal birth outcome.<sup>3,4</sup> 2. It is truly reassuring for women with IBD that IBD is not incompatible with maternity. They could conceive safely, provided that they are in the remission stage, and under the close supervision of multidisciplinary team involving not only an obstetrician and gastroenterologist, but also a pediatrician to monitor the disease activity, pregnancy outcome, and the health status of the newborn.

Mahmood D. Al-Mendalawi
Department of Pediatrics
Al-Kindy College of Medicine
Baghdad University
Baghdad, Iraq

#### Reply from the Author

No reply was received from the Author.

### References

- Saleh A, Badroun HI, Al-Khaldi LW, Khalifa HM, Waggass MA. Pregnancy outcome in women with inflammatory bowel disease. *Saudi Med J* 2010; 31: 1174-1176.
- Raatikainen K, Mustonen J, Pajala MO, Heikkinen M, Heinonen S. The effects of pre- and post-pregnancy inflammatory bowel disease diagnosis on birth outcomes. *Aliment Pharmacol Ther* 2011; 33: 333-339.
- Mahadevan U, Sandborn WJ, Li DK, Hakimian S, Kane S, Corley DA. Pregnancy outcomes in women with inflammatory bowel disease: a large community-based study from Northern California. *Gastroenterology* 2007; 133: 1106-1112.
- Molnár T, Farkas K, Nagy F, Lakatos PL, Miheller P, Nyári T, et al. Pregnancy outcome in patients with inflammatory bowel disease according to the activity of the disease and the medical treatment: a case-control study. *Scand J Gastroenterol* 2010; 45: 1302-1306.

#### Related topics

Shuaib AA, Frass KA, Al-Harazi AH, Ghanem NS. Pregnancy outcomes of mothers aged 17 years or less. *Saudi Med J* 2011; 32: 166-170.

Bhatti KA, Babay ZH, Neyazi SM. Ruptured spontaneous heterotopic pregnancy. *Saudi Med J* 2010; 31: 445-447.

Mansouri HA. Central nervous system anomalies diagnosed antenatally and post-delivery management. *Saudi Med J* 2010; 31: 257-261.

Abu-Skeen IA, Mohamed AA, Moustafa NN, Badawy ME. Factor V Leiden and prothrombin G20210A gene mutations in women with a history of thrombosis during pregnancy. *Relation to pregnancy outcomes for mother and fetus. Saudi Med J* 2010; 31: 123-129.