



## WHO URGES ACTION AGAINST HIV DRUG RESISTANCE THREAT

20 JULY 2017 | GENEVA - WHO alerts countries to the increasing trend of resistance to HIV drugs detailed in a report based on national surveys conducted in several countries. The Organization warns that this growing threat could undermine global progress in treating and preventing HIV infection if early and effective action is not taken.

The WHO HIV drug resistance report 2017 shows that in 6 of the 11 countries surveyed in Africa, Asia and Latin America, over 10% of people starting antiretroviral therapy had a strain of HIV that was resistant to some of the most widely used HIV medicines. Once the threshold of 10% has been reached, WHO recommends those countries urgently review their HIV treatment programmes.

### HIV drug resistance report 2017

“Antimicrobial drug resistance is a growing challenge to global health and sustainable development,” said Dr Tedros Adhanom Ghebreyesus, WHO Director-General. “We need to proactively address the rising levels of resistance to HIV drugs if we are to achieve the global target of ending AIDS by 2030.”

HIV drug resistance develops when people do not adhere to a prescribed treatment plan, often because they do not have consistent access to quality HIV treatment and care. Individuals with HIV drug resistance will start to fail therapy and may also transmit drug-resistant viruses to others. The level of HIV in their blood will increase, unless they change to a different treatment regimen, which could be more expensive – and, in many countries, still harder to obtain.

Of the 36.7 million people living with HIV worldwide, 19.5 million people were accessing antiretroviral therapy in 2016. The majority of these people are doing well, with treatment proving highly effective in suppressing the HIV virus. But a growing number are experiencing the consequences of drug resistance.

WHO is therefore issuing new guidelines to help countries address HIV drug resistance. These recommend that countries monitor the quality of their treatment programmes and take action as soon as treatment failure is detected.

“We need to ensure that people who start treatment can stay on effective treatment, to prevent the emergence of HIV drug resistance,” said Dr Gottfried Hirnschall, Director of WHO’s HIV Department and Global Hepatitis Programme. “When levels of HIV drug resistance become high we recommend that countries shift to an alternative first-line therapy for those who are starting treatment.”

Increasing HIV drug resistance trends could lead to more infections and deaths. Mathematical modelling shows an additional 135 000 deaths and 105 000 new infections could follow in the next five years if no action is taken, and HIV treatment costs could increase by an additional US\$ 650 million during this time.

Tackling HIV drug resistance will require the active involvement of a broad range of partners. A new five-year Global Action Plan calls on all countries and partners to join efforts to prevent, monitor and respond to HIV drug resistance and to protect the ongoing progress towards the Sustainable Development Goal of ending the AIDS epidemic by 2030. In addition, WHO has developed new tools to help countries monitor HIV drug resistance, improve the quality of treatment programmes and transition to new HIV treatments, if needed.

The WHO HIV drug resistance report 2017 was co-authored by the Global Fund to Fight AIDS, Tuberculosis and Malaria, and the Centers for Disease Control and Prevention, USA.

“This new report shows a worrying picture of increasing levels of HIV drug resistance and, if unchecked, it will be a major risk to program impact,” said Dr Marijke Wijnroks, Interim Executive Director of the Global Fund. “We strongly recommend implementing WHO recommendations for early warning indicators and HIV drug resistance surveys in every national plan for antiretroviral therapy, and to consider funding them through Global Fund grants or reprogramming.”

Dr Shannon Hader, Director of CDC’s Division of Global HIV and Tuberculosis, US Centers for Disease Control and Prevention, added: “The new report pulls together key HIV drug resistance survey findings from across the globe that, taken together with other national-level data, confirm we must be forward-thinking in our efforts to combat resistance: scaling up viral load testing, improving the quality of treatment programs, and transitioning to new drugs like dolutegravir.”

Dr. Hader continued, stating that “Overall high rates of viral suppression across three recent national Population-based HIV Impact Assessments showed that present first-line regimens remain largely effective. However, special attention to populations at risk for higher resistance, such as pediatrics, adolescents, pregnant women and key populations, will be critical to target more urgent interventions. We call on the global community for continued vigilance and responsiveness.”

Available from: <http://www.who.int/mediacentre/news/releases/2017/hiv-drug-resistance/en/>

---

## BABIES AND MOTHERS WORLDWIDE FAILED BY LACK OF INVESTMENT IN BREASTFEEDING

1 AUGUST 2017 | No country in the world fully meets recommended standards for breastfeeding, according to a new report by UNICEF and WHO in collaboration with the Global Breastfeeding Collective, a new initiative to increase global breastfeeding rates.

### Global Breastfeeding Collective

The Global Breastfeeding Scorecard, which evaluated 194 nations, found that only 40 per cent of children younger than six months are breastfed exclusively (given nothing but breastmilk) and only 23 countries have exclusive breastfeeding rates above 60 per cent.

Evidence shows that breastfeeding has cognitive and health benefits for both infants and their mothers. It is especially critical during the first six months of life, helping prevent diarrhoea and pneumonia, two major causes of death in infants. Mothers who breastfeed have a reduced risk of ovarian and breast cancer, two leading causes of death among women.